arXiv:1304.5703v1 [cond-mat.soft] 21 Apr 2013

Dipolar Self Assembly in Coherently Fluctuating Fields

Igor M. Kulić[1]* and Miodrag L. Kulić[2]†
[1] *CNRS, Institute Charles Sadron, 23 rue du Loess BP 84047, 67034 Strasbourg, France and*
[2] *Institute for Theoretical Physics, Goethe-University D-60438 Frankfurt am Main, Germany*

(Dated: April 23, 2013)

When two polarizable objects interact in spatially randomly fluctuating fields, we speak of Van der Waals-like forces. But what happens if the random excitation becomes spatially coherent? We show that such a *spatially coherent fluctuation interaction* (scFI) displays remarkable physical features distinguishing it from its sister, the incoherent fluctuation interaction (icFI). In particular, we investigate how scFI gives rise to complex, hierarchically structured, self-assembled materials, in contrast to the usual phase-separated droplets generated by icFI. We discuss a recent example of scFI, implemented with superparamagnetic colloids in triaxial magnetic fields by Martin et al. and explore the main steps in the hierarchical formation process of coherent colloid self assemblies, from chains, through membranes, to large-scale foamy superstructures of the material. We predict that such coherent "dipolar doughs" exert positive pressure, can easily swell against gravity on cm-scales and we derive their equation of state.

PACS numbers: 82.70.Dd, 81.16.Dn, 82.70.Rr

The basic motif behind the formation of most solids and liquids is the interplay of a long-range attractive and a short-range repulsive force. As happens often in physical systems, simply reversing the signs of the interactions is a real game changer. A competing short-range attractive and long-range repulsive force can give rise to profoundly different, complex, sponge-like structures ranging from gyroid phases in block copolymers [1], labyrinthine phases in ferrofluids [2] to nuclear pasta phases in gravity-collapsed neutron stars [3]. By imposing spatially coherent conically rotating (triaxial) magnetic fields Martin and coworkers have discovered [4, 5] that superparamagnetic *Ni*- colloids, placed in a finite-volume container, self-assemble into unusual foam-like structures, displaying a local cohesive force on small scales but with an overall large-scale tendency to repel and rise against gravity, much like a swelling dough (cf. Fig.4a below). Owing to the complexity of the spatially coherent oscillating field-induced magnetic interaction and its inherent anisotropic dipolar nature, the structures formed on the smallest scales unexpectedly resemble mono-layered colloidal *membranes* [4–6] - in sharp contrast to the chains predominating in the usual time-constant fields. After a continued growth phase the colloidal membranes begin to merge into larger-scale foam-like superstructures spanning the whole sample container. What Martin et al. have discovered in their seminal work [4, 5], is a particular instance of a *spatially coherent fluctuation interaction* (scFI): a spatially homogeneous dipole-exciting field with certain statistical properties (its square is statistically isotropic) varies in time and gives rise to an effective interaction (cf. Fig.1) of amazing complexity with a significant potential for micro- and nano-scale self-assembly.

This letter aims at bringing order into our theoretical understanding of scFI by outlining a certain simplicity underlying its seeming complexity. We will explore how scFI's intrinsically strong many-body interactions drive the formation of chains and membranes and finally how complex membrane-membrane pair interactions give rise to swelling "dipolar doughs", whose equation of state we derive at the end of the letter.

1.Coherent vs. Incoherent Fluctuation Interaction. How can the scFI (cf. Fig.1) generate such complex structures while its sister - the *incoherent fluctuation interaction* (icFI)-merely forms phase-separated lumps or droplets of matter within a two phase system [7]? To answer this question we consider dipolar magnetic (dielectric) particles without permanent moments [8], such as colloids, which are placed in a temporally fluctuating magnetic (electric) field $\mathbf{B}_{0,i} = \mu_0 \mathbf{H}_{0,i}(t)$, which changes on a typical timescale t_ω. The latter fulfills the condition $\tau_M < t_\omega < \tau_\eta$ with τ_M the dipolar relaxation time and τ_η the characteristic time for the colloid's motion in the surrounding viscous fluid. Under these conditions the beads' magnetization is equilibrated, while their centers of mass respond much more slowly and feel a net time-averaged force. The free energy functional $\mathcal{F}$ for interacting beads with magnetization $\mathbf{M}_i$ in imposed fields $\mathbf{H}_{0,i}$ is then given by [9]

$$\frac{1}{\mu_0}\mathcal{F}(\mathbf{M}_i;\mathbf{H}_{i,0}) = \sum_i \left(\frac{1}{2}\mathbf{M}_i\hat{\chi}_{b,i}^{-1}\mathbf{M}_i - \mathbf{M}_i\mathbf{H}_{0,i}\right)V_i \tag{1}$$

$$+\frac{1}{2}\sum_{i,j\neq i}\mathbf{M}_i\hat{T}_{ij}\mathbf{M}_j V_i,$$

where the summation goes over N beads, μ_0 is the vacuum permeability, V_i is the volume of the i-th bead and $\hat{\chi}_b$ is the *bead susceptibility* tensor. For an ellipsoidal colloid (or colloid aggregate), one has $\hat{\chi}_b = (1+\hat{L}_b\chi_{mb})^{-1}\chi_{mb}$, where χ_{mb} is the colloid's *material susceptibility* - which can take very large values in multidomain (or superparamagnetic) beads - and $\hat{L}_b$ is its demagnetization tensor. The first term in Eq.1 comes from the ferromagnetic domain entropy within beads, while the bead-bead magnetic interaction is contained in the third term [10]. Here we keep only the magnetic dipole-dipole interaction [4, 5], given by $\hat{T}_{ij} = \varphi_{ij}\hat{t}(\mathbf{b}_{ij})$, with $\varphi_{ij} \equiv \varphi(\mathbf{R}_i, \mathbf{R}_j) = V_j/4\pi\left|\mathbf{R}_{ij}\right|^3$, $\mathbf{R}_{ij} \equiv \mathbf{R}_i - \mathbf{R}_j$ and the tensor $\hat{t}(\mathbf{b}_{ij}) = \hat{1} - 3\mathbf{b}_{ij} \otimes \mathbf{b}_{ij}$ (dyadic product) with the "bonding" unit vector $\mathbf{b}_{ij} = \mathbf{R}_{ij}/\left|\mathbf{R}_{ij}\right|$. After minimizing $\mathcal{F}$ with respect to $\mathbf{M}_i$ for identical beads ($V_{\mathbf{i}} = V_b$), followed by time-averaging over a spatially coherent excitation, $\mathbf{H}_{0,i} = \mathbf{H}_0$, $\overline{H_{i,0\alpha}H_{j,0\beta}} = \delta_{\alpha\beta}H_0^2$ ($i, j = 1, .., N$; $\alpha, \beta = x, y, z$), one obtains a physically transparent expression for the effective energy in the scFI case

$$\bar{\mathcal{F}}_{scDI}\left(\mathbf{H}_0, \{\mathbf{R}_i\}\right) = -\frac{\mu_0}{2}H_0^2 V_b \sum_{i,j} Tr\left\{\hat{\chi}_{eff,ij}\right\} \qquad (2)$$

where $Tr\left\{\hat{\chi}_{eff,ij}\right\} = \chi^{xx}_{eff,ij} + \chi^{yy}_{eff,ij} + \chi^{zz}_{eff,ij}$ is the spatial trace of the (non-local) *effective susceptibility* $\hat{\chi}_{eff,ij} = \chi_b \left\langle \mathbf{R}_i \right| (\hat{1} + \chi_b\hat{T})^{-1} \left| \mathbf{R}_j \right\rangle$ with $\left\langle \mathbf{R}_i \right| \hat{T} \left| \mathbf{R}_j \right\rangle \equiv \hat{T}_{ij}$. As $\hat{\chi}_{eff,ij}$ describes the coupling of the i-th and j-th bead *in presence of all other beads*, it gives rise to many-body effects. These are crucial for the hierarchical structuring of colloids, as confirmed in numerical simulations [4, 5] and experiments [4–6], where dimers, molecules, chains, branched chains, membranes and doughs are formed, instead of droplets and close-packed 3D crystal structures. The elegant "trace-formula" Eq.2 forms the basis for our further study of scFI systems. Note here that, in contrast to scFI, in standard incoherently excited FI (icFI) systems, with $\overline{H_{i,0\alpha}H_{j,0\beta}} = \delta_{ij}\delta_{\alpha\beta}H_0^2$ (note the δ_{ij}), the summation of $Tr\left\{\hat{\chi}_{eff,ij}\right\}$ in $\bar{\mathcal{F}}$ includes only $i = j$ terms, giving rise to the usual Van der Waals(VdW) -like forces, preferring the formation of simple droplets only [7]. In the following, we study first the bead-bead interaction, then the formation of chains and membranes, and finally the assembly of membranes in the form of $3D$ doughs.

2. Dimer formation. In a first step, let us consider a very dilute system, where the *pairwise* bead-bead interactions should (naively) dominate in $\bar{\mathcal{F}}_{scDI}$. For two spherical beads (with indices 1,2), we have $\hat{L}_{\alpha\beta} = (1/3)\delta_{\alpha\beta}$, $(\hat{\chi}_b)_{\alpha\beta} = \chi_b\delta_{\alpha\beta}$ (with $\chi_b \leq 3$) and Eq.(2), gives $Tr\left\{\hat{\chi}^{(b)}_{eff,12}\right\} = 3\chi_b(1 - \chi_b\varphi_b)/(1 + \chi_b\varphi_b)(1 - 2\chi_b\varphi_b)$, where $\varphi_b = V_b/4\pi\left|\mathbf{R}_{12}\right|^3$. This means that the bead-bead interaction is attractive for $\chi_b > 0$. For large distances $\left|\mathbf{R}_{12}\right| \gg V_b^{1/3}$, the interaction is $\bar{\mathcal{F}}^{(bb)}_{12} = -(3/8\pi^2)(V_b\chi_b)^3H_0^2\left|\mathbf{R}_{12}\right|^{-6}$ giving rise to an isotropic and short-range attractive VdW-like attractive force $\mathbf{F_{12}} = -(\partial\bar{\mathcal{F}}^{(bb)}_{12}/\partial\mathbf{R}_1) \propto \left|\mathbf{R}_{12}\right|^{-7}$. This interesting two-body result was first obtained in the seminal papers [4, 5] and confirmed experimentally [6].

3.Many-body effects. At first glance, the isotropic VdW-like two-body interaction $\propto 1/\left|\mathbf{R}_{12}\right|^{-6}$ appears to favor droplet-like, bulk structures. However, this contradicts numerical simulations by Martin et al. [4, 5] and experiments [4–6], which show a clear tendency for chain and membrane formation. What is the microscopic origin of these complex structures in scFI systems? The reason hides in the specificity and the strength of the many-body interactions, which mark the real difference between scFI and icFI systems. In a typical icFI system, the 3-body interaction is described by an Axilrod-Teller type 3-body potential $\bar{\mathcal{F}}^{(123)}_{A-T} = C_{123}/\left|\mathbf{R}_{12}\right|^3\left|\mathbf{R}_{23}\right|^3\left|\mathbf{R}_{31}\right|^3$, with $C_{123} \propto \chi_b^3\left(1 + \cos\theta_1\cos\theta_2\cos\theta_3\right)$ [11]. Due to its weaker $\sim \chi^3R^{-9}$ scaling, it is typically small and overridden by the 2-body VdW-like interaction $\sim \chi^2R^{-6}$ [12], giving rise to close packed droplets in icFI systems. In sharp contrast, the 3-body interaction $\bar{\mathcal{F}}^{(123)}_{scFI}$ in scFI systems scales very differently with distance, as seen from a $O\left(\chi_b^3\varphi_b^3\right)$ expansion of Eq.(2) [13]:

$$\bar{\mathcal{F}}^{(123)}_{scFI} = -\frac{3}{32\pi^2}\mu_0H_0^2\chi_b^3V_b^3\frac{3\cos^2\theta_1 - 1}{\left|\mathbf{R}_{12}\right|^3\left|\mathbf{R}_{13}\right|^3} + perm. \qquad (3)$$

where we sum over all 3-index permutations (for angles $\theta_{1,2,3}$ cf. Fig.1d). Remarkably, $\bar{\mathcal{F}}^{(123)}_{scFI}$ is of the same order in R and χ_B as the 2-body interaction $\bar{\mathcal{F}}^{(12)}_{scFI} \sim -\chi_b^3/\left|\mathbf{R}_{12}\right|^6$. In fact, the 2-body interaction is formally *contained* in $\bar{\mathcal{F}}^{(123)}_{scFI}$ (indices 1≠2=3) showing us the pitfall in the "Dimer formation" section: For *scFI* the 2-body interactions are physically *inseparable* from the 3-body ones - at any χ_b! Interestingly, $\bar{\mathcal{F}}^{(123)}_{scFI}$ has a simple angular dependence explaining scFI's tendency to destroy 3D bulk structures: the $-\cos^2\theta_i$ term favors $\theta_i = 0$ or π - i.e. the colloidal chains found in the experiments



FIG. 1: a) The standard incoherent (icFI) and the b) spatially coherent fluctuation interaction (scFI) are both induced by field fluctuations - but with different spatial correlations. c)+d) While the 2-body forces appear superficially similar in both cases, the 3-body forces have different angular character and are longer ranged for scFI, Eq.(3).



FIG. 2: Left panel: scFI leads to the formation of colloidal clusters with random orientations growing over time [4, 5]. At the early stages, smaller clusters are born as linear chains and transformed into membranes beyond a critical size $N_c\,(\chi)$. Right panel: Phase diagram of clusters of size N and susceptibility χ.

[4–6].

4. Chains and membranes. From the 3-body forces of Eq.3 we understand now intuitively why chains form initially. To capture their transition to membranes, higher $O\left(\chi_b^4\varphi_b^4\right)$ terms beyond Eq.3 are necessary. It is however conceptually more instructive to take a more macroscopic approach, where chains/membranes are replaced by prolate/oblate ellipsoids. Here, $\hat{\chi}_{eff,ij}$ in Eq.2 is replaced by its shape-dependent continuum limit $\hat{\chi}^{(L)}$, which is spatially homogeneous inside ellipsoidal samples with dimensions $a = b \neq c$ and volume $V = (4\pi/3)a^2c$. The free-energy $\bar{\mathcal{F}}_{self}$ and $\hat{\chi}^{(L)}$ are in this case given by

$$\bar{\mathcal{F}}_{self}(\mathbf{H}_0, \hat{L}) = -\frac{1}{2}\mu_0 H_0^2 V Tr\left\{\hat{\chi}^{(L)}\right\} \qquad (4)$$
$$\hat{\chi}^{(L)} = \chi(1+\hat{L}\chi)^{-1},$$

where $\hat{L}$ is now the demagnetization tensor of the composite ellipsoidal structure, with $Tr\hat{L} = L_a + L_b + L_c = 1$ and L_i its eigenvalues [9]. Here, $\chi \approx \chi_b f(1-\chi_b f/3)^{-1}$ is the *composite material susceptibility,* which is due to local field effects in the dense bead aggregate, with $f(<1)$ the volume fraction of beads within the membrane/chain.

In the continuum limit, when the number of beads is large ($N \gg 1$), chains can be considered as *extreme prolate ellipsoids* with a long half-axis $c \sim ND$, a short half-axis $a \sim D$ and the demagnetization factors $L_a = L_b \gg L_c \equiv L^{ch} \sim (\ln N)N^{-2}$. In that case one has $Tr\left\{\hat{\chi}^{ch}\right\} = \chi(1+L^{ch}\chi)^{-1} + 2\chi[1+(1-L^{ch})\chi/2]^{-1}$ in Eq.4. Membranes, on the other hand, can be considered as *extreme oblate ellipsoids* with two identical half-axes $a = b \sim \sqrt{N}D$, $c \sim D$ and $L_c \gg L_a = L_b \equiv L^m \sim N^{-1/2}$. In this case one has $Tr\left\{\hat{\chi}^m\right\} = 2\chi(1+L^m\chi)^{-1} + \chi[1+(1-2L^m)\chi]^{-1}$ in Eq.4. In both cases, for chains and membranes, $\bar{\mathcal{F}}_{self}$ is dominated by the smallest demagnetization factors $L^{m/ch} \to 0$, in the limit $N \to \infty$. However, since for membranes two demagnetization factors vanish, while for chains only one vanishes, the energy of a membrane is always smaller than that of the chain for $N \to \infty$, i.e. $\bar{\mathcal{F}}^m < \bar{\mathcal{F}}^{ch}$. For finite N, the *chain-membrane transition* is reached on the line $N = N_c(\chi)$, where $\bar{\mathcal{F}}^m(N_c) = \bar{\mathcal{F}}^{ch}(N_c)$. As seen from Fig.2, N_c grows with the composite susceptibility χ. For $\chi \approx 3$ (for "Dynabeads" with $\chi_b \approx 1$ [6]), we estimate $N_c \approx 10-15$ - close to the experimental value $N_c \approx 10$ for the chain-membrane transition [6]. The agreement suggests that the continuum approach quantitatively captures the behavior for finite N.

5. Interaction of membranes - Once they emerge, what is the fate of the membranes as they continue growing? How do they interact and mutually order during growth? To answer these questions, we need the *2-membrane interaction* for arbitrary membrane orientations $\mathbf{n}_{1,2}$ and anisotropic susceptibilities (of thin oblates) $\hat{\chi}_i^{(L)} = \chi(1+\hat{L}_i\chi)^{-1}$. Up to linear order in φ_m, one expands $\hat{\chi}_{eff,1} \approx \hat{\chi}_1^{(L)}(1-\varphi_m\ \hat{t}(\mathbf{b}_{12})\hat{\chi}_{\mathbf{2}}^{(L)})$ (and $\hat{\chi}_{eff,\mathbf{2}}$ for 1→2). For identical, distant membranes ($V_{m,1/2} = V_m, |\mathbf{R}_{12}| \gg V_m^{1/3}$), the $O\left(\varphi_m\right)$ interaction energy in Eq.(2) reads

$$\bar{\mathcal{F}}_{int} = \frac{3\mu_0 H_0^2 V_m^2 (1-\gamma)\chi_{\max}^2}{16\pi\left|\mathbf{R}_{12}\right|^3} \left(c_1^2 + c_2^2 + \frac{1-\gamma}{3}c_3^2 - (1-\gamma)c_1c_2c_3 - \frac{2}{3}\right) \qquad (5)$$

with $\chi_{\min\,/\,\max}$ the minimal/maximal eigenvalue of the membrane susceptibility tensor $\hat{\chi}^{(L)}$, $\gamma = \chi_{\min}/\chi_{\max}$ and geometric factors $c_1 = \mathbf{n}_1\cdot\mathbf{b}_{12}$, $c_2 = \mathbf{n}_2\cdot\mathbf{b}_{12}$, $c_3 = \mathbf{n}_1\cdot\mathbf{n}_2$. This 2-membrane interaction is angle dependent and repulsive in many configurations - see Fig.3. Notably, for fixed $\left|\mathbf{R}_{12}\right|, \bar{\mathcal{F}}_{int}$ becomes minimal for the orthogonally "*twisted*" membrane orientation with $\mathbf{n}_1 \perp \mathbf{n}_2$, $\mathbf{n}_1 \perp \mathbf{b}_{12}$ and $\mathbf{n}_2 \perp \mathbf{b}_{12}$ ($c_{1/2/3} = 0$). The "twisted" membranes attract each other since $\bar{\mathcal{F}}_{int}^{(tw)} < 0$ (up to the point of mutual contact), as in the "*coplanar*" case, yet the "*twisted*" configuration has lower energy. This interesting result should affect the kinetics of membrane formation: If two distant membranes start growing within a large L they will rotate to a 90° position before touching. Therefore, in finite Ls, some type of glassy state in their orientation may be kinetically favored. In other relevant configurations, such as the "*top*", with two out of plane parallel membranes ($c_{1/2/3} = 1$) or the "*generic*" one (cf. Fig.3), the interaction is repulsive with $0 < \bar{\mathcal{F}}_{int}^{(gen)} < \bar{\mathcal{F}}_{int}^{(top)}$.

6.Formation of doughs and equation of state. Simulations and experiments [4, 5] provide empirical evidence for a hollow dough-like superstructure in finite Ls (cf. Fig 4a). What is the physical mechanism behind such "dipolar dough" formation? We have seen above that large aggregates prefer to form membranes, and that these membranes mutually interact. Specifically, when two distant membranes are stacked over each other, they repel



FIG. 3: The scFI is unexpectedly complex in various geometries. a) The 2-bead interaction is simple in comparison, always attractive and $\propto -R^{-6}$ in leading order. b) The bead-membrane interaction is longer ranged and anisotropic in relative coordinates, changing scaling to $\propto \pm R^{-3}$. The interaction flips sign from attractive to repulsive on a cone with opening angle $\theta_{crit} \approx 54.7°$ [13] (c). The 2-membrane interaction is complex, attractive or repulsive depending on orientation, cf. Eq.5, with a ground state in the "twisted" configuration.



FIG. 4: (a) The large scale structure of a dipolar dough (from [4, 5]) in experiment (left) and in simulation (right). (b) The theoretical 3D shelf-model for dough's structure.

($\bar{\mathcal{F}}_{int}^{(top)} > 0$). In the opposite limit (contact distance), a simple estimate implies their preference to split as well [14]. It is this remarkable reluctance of membranes to mutually stack that in fact sets the microscopic structure of the "dough". The dough is formed out of the thinnest possible membrane patches, whose *thickness* is collapsed onto the smallest available physical scale - the colloid size D. The characteristic lateral size a_M of these membrane patches ,on the other hand, is set by the bead volume fraction in the container $f_V = V_b^{tot}/V \ll 1$, with V_b^{tot} the total volume of all beads and V the container's volume. By assuming a cubic shelf structure as an ansatz (cf. Fig 4b) one obtains a patch size $a_M = 3D/f_V$.

Since the membranes tend to grow, yet repel on the average, the container walls will feel a net pressure $p(= -(\partial\bar{\mathcal{F}}/\partial V)) > 0$ [15], where $\bar{\mathcal{F}} = \bar{\mathcal{F}}_{self} + \bar{\mathcal{F}}_{int}$ has contributions from both the membrane self-energy $\bar{\mathcal{F}}_{self}$ (from Eq.4) and their pairwise interactions $\bar{\mathcal{F}}_{int}$ (Eq.(5)). Since both attractive and repulsive membrane interactions are present, it is a difficult endeavor to find the exact equilibrium structure, even in numerical calculations. However, using the simple shelf-like cubic lattice ansatz with the mesh-size a_M (Fig.4), we can qualitatively mimic the dough structure [4, 5]. For this regular lattice, we can sum the pairwise membrane interactions and obtain the total interaction energy (for $\chi \gg 1$) $\bar{\mathcal{F}}_{int} \approx (S/8\pi)V_b^{tot} f_V \chi^2 p_0$, with $p_0 = B_0^2/2\mu_0$ and the lattice interaction constant [13] $S = \sum_{i\neq 1}\left(c_1^2 + c_2^2 + \frac{1}{3}c_3^2 - c_1c_2c_3 - \frac{2}{3}\right)|a_M/\mathbf{R}_{1i}|^3 \approx 10$ obtained numerically from Eq.5.

The self energies of finite membranes depend on their demagnetization factor L^m (see discussion above), which for the shelf structure is given by f_V as $L^m \approx \alpha_m f_V$ for $f_V \ll 1$ with $\alpha_m \approx 1/4$. For $\chi \gg 1$ and $L^m\chi \ll 1$ (dilute limit) the self-energy is given then by $\bar{\mathcal{F}}_{self} \approx const + 0.5\chi^2 f_V p_0 V_b^{tot}$. The corresponding contributions to the pressure are $p_{self} \approx 0.5 f_V^2 \chi^2 p_0$ and $p_{int} \approx 0.4 f_V^2 \chi^2 p_0$ respectively, thus giving $p \approx f_V^2 \chi^2 p_0$. The compressibility of the dough is $\kappa = 1/2p$. For $f_V \approx 5 \cdot 10^{-2}$ and $B_0 = 10\ mT$, we obtain $p \sim 10\ Pa$ and a large $\kappa \sim 0.1\ Pa^{-1}$. Since $\kappa \propto B_0^{-2}$ the compressibility is sensitive to the strength of excitation B_0. Finally, in a gravitational field the *maximal height* $h_{\max}$ of the dough is reached for $p \approx p_{gr}$, with the gravitational pressure $p_{gr} = \Delta\rho g h f_V$. For water immersed Ni-beads [4, 5] one has the density contrast $\Delta\rho \simeq 8 \cdot 10^3 kg/m^3$ and $\chi \sim 10$ in a densely packed membrane. For $f_V \sim 5 \cdot 10^{-2}$, $B_0 = 20\ mT$ and $\chi = 10$ we predict that the dough rises significantly, with $h^0 \sim 1\ cm$ [16].

In conclusion, we have studied, in the framework of the effective nonlocal susceptibility $\hat{\chi}_{ij}$, the formation of hierarchical superstructures in dipolar colloids driven by the spatially coherent fluctuation interaction (scFI). The pronounced many-body interactions (contained in $\hat{\chi}_{ij}$) give rise to a growth of anisotropic assemblies - chains, then membranes once a critical cluster size $N_c(\chi)$ is reached. In a spherical container of finite size, smaller membrane patches of size $\sim D/f_V$ are formed, which,contrary to the case of attracting beads, repel on average, thus giving rise to "dipolar dough" structures. The dough exerts a positive pressure on the walls of the container due to the tendency of membranes to increase their surface areas as well as their mutual repulsion. The "dipolar dough" represents an intriguing state of colloidal matter, formed by a subtle interplay of an attractive local interaction and a net repulsive longer range force. Remarkably, both types of forces are born out of a single, conceptually simple interaction - the scFI (Eq.2). The equation of state $p \sim f_V^2 \chi^2 B_0^2$ gives rise to a large compressibility, highly sensitive to changes of magnetic field, opening doors for applications. Dielectric scFI analogues, like light-illuminated plasmonic nanoparticle systems, are obvious next challenges. Here, dynamic susceptibility effects near resonance frequencies should add new physics needed to understand the shape-resonant self-assemblies found in Ref.[17]. The complexity of the quasi-static scFI considered here makes us wonder about the laws of the nano-cosmos emerging once dynamic effects are switched on.

Acknowledgements. We thank A. Johner, H. Mohrbach and M. Greenall for discussions and useful comments.

---

[*] Electronic address: kulic@unistra.fr
[†] Electronic address: kulic@th.physik.uni-frankfurt.de

[1] I. W. Hamley, The Physics of Block Copolymers , Oxford Science Publications, (1999)

[2] A. Tsebers, M. Maiorov. Magnetohydrodynamics, vol. 16, 21–27 (1980); Rosensweig R E, Zahn M and Shumovich R J. Magn. Magn. Mater. 39, 127-32 (1983)

[3] D. G. Ravenhall, C. J. Pethick, and J. R. Wilson, Phys. Rev.Lett. 50, 2066 (1983); M. Hashimoto, H. Seki, and M. Yamada, Prog. Theor. Phys. 71, 320 (1984).

[4] J. E. Martin, R. A. Anderson, R. L. Williamson, J. Chem. Phys. **118**, 1557 (2003)

[5] J. E. Martin, E. Venturini, G. L. Gulley, J. Williamson, Phys. Rev. **E 69**, 021508-1 (2004)

[6] N. Osterman, I. Poberaj, J. Donikar, D. Frenkel, P.Ziherl, D. Babić, Phys. Rev. Lett. **103**, 228301 (2009)

[7] V.A. Parsegian, Van Der Waals Forces, Cambridge University Press (2006)

[8] We adopt the magnetic notation but all results apply in the electric formalism (paramagnets become dielectrics).

[9] L. D. Landau, E. M. Lifshitz, Electrodyanamics of Continuous Media, Oxford: Pergamon Press (1989)

[10] Due to the large moments $\mathbf{m}_i = \mathbf{M}_i V_i$ of the beads with diameters $D > 1\ \mu m$, $\mathcal{F} \gg k_B T$ making additional contributions of a configurational entropy negligible.

[11] B. M. Axilrod, E. Teller, J. Chem. Phys. **11**, 299 (1943); Yu. S. Barash, V. L. Ginzburg, Sov. Phys. Uspekhi, **143**, 345 (1984)

[12] The χ^2 scaling is valid for equilibrium conditions, where a detailed ballance between the moments and (thermal or quantum) bath hold. The scaling for icFI switches to $\propto \chi^3$ if the fluctuating field is exogeneous (externally set) and non-equilibrium as in the present scFI case [13].

[13] I. M. Kulić and M. L. Kulić, to be published elsewhere

[14] When a thick membrane (thickness $2D$, radius R and volume $2V_m$) is cut into two parallel membranes (thickness D and radius R each) and separated to infinite distance there is a gain in the energy $\Delta\mathcal{F} = 2\mathcal{F}_{1m} - \mathcal{F}_{2m} \approx -V_{2m} L \chi^2 (1 - (1+\chi)^{-2}) < 0$ for $L\chi \ll 1$, where $L \propto D^{3/2} V_{2m}^{-1/2}$. Physically, the second membrane lying above the first one is repelled to increase the local fields with respect to the thicker membrane case [13].

[15] In general p is container shape dependent and we expect $p^{obl} > p^{sph} > p^{pro} > 0$ for the oblate, spherical and prolate containers respectively.

[16] More precisely, due to gravity, f_V and p vary with h, i.e. $p(f_V) = p_{\max}(1 - h/2h^0)^2$ giving the maximal height $h_{\max} = 2h^0 \sim 2\ cm$.

[17] K.J. Stamplecoskie & J.C. Scaiano, JACS 132, 1825 (2010)